P130a   BISTRO Project Status (3)

Tetsuo Hasegawa[1], Ray Furuya[2], Doris Arzoumanian[3], Yasuo Doi[4], Saeko Hayashi[1], Charles Hull[1], Tsuyoshi Inoue[3], Shu-ichiro Inutsuka[3], Kazunari Iwasaki[5], Yoshihiro Kanamori[4], Akimasa Kataoka[1], Koji Kawabata[6], Masato Kobayashi[3], Takayoshi Kusune[7], Jungmi Kwon[8], Masafumi Matsumura[9], Tetsuya Nagata[10], Fumitaka Nakamura[1], Hiroyuki Nakanishi[11], Nagayoshi Ohashi[1], Takashi Onaka[4], Katherine Pattle[1,15], Tae-Soo Pyo[1], Hiro Saito[12], Masumichi Seta[13], Hiroko Shinnaga[11], Motohide Tamura[4,14], Kohji Tomisaka[1], Yusuke Tsukamoto[11], Tetsuya Zenko[10], Derek Ward-Thompson[15] and BISTRO Consortium ([1]NAOJ, [2]Tokushima U., [3]Nagoya U., [4]U. Tokyo, [5]Doshisha U., [6]Hiroshima U., [7]Nagoya City U., [8]ISAS, [9]Kagawa U., [10]Kyoto U., [11]Kagoshima U., [12]Tsukuba U., [13]Kwansai U., [14]Astro Biology Center, [15]U. of Central Lancashire)

BISTRO (B-field In STar forming Region Observations) is an international research project to make sub-millimeter linear polarization images of nearby star forming regions as one of the EAO/JCMT Large Programs that involves over a hundred researchers in Canada, China, Japan, Korea, Taiwan, UK, and the East Asian Observatory. The data taking is almost complete for the BISTRO sources (Orion A/B, rho Oph, Serpens Main, Perseus B1, IC5146, NGC1333, Taurus B211/213, L1495), and two papers have been published with several more submitted or in preparation in the international consortium. The JCMT SCUBA-2/POL-2 and the data reduction software achieve about an order of magnitude improvement of the dynamic range of the polarization images. This brings a breakthrough to our understanding to the role of magnetic field in star formation.